

National Park Service  
U.S. Department of the Interior

Natural Resource Program Center



# Methods for Monitoring Landbirds

*A Review Commissioned by Seattle City Light's Wildlife  
Research Advisory Committee (2000)*

Natural Resource Report NPS/NCCN/NRR—2009/074



**ON THE COVER**

Common yellowthroat

Photograph by Harold Christianson

---

**Methods for Monitoring Landbirds**  
*A Review Commissioned by Seattle City Light's Wildlife  
Research Advisory Committee (2000)*

Natural Resource Report NPS/NCCN/NRR—2009/074

Rodney B. Siegel  
The Institute for Bird Populations  
P.O. Box 1346  
Point Reyes Station, CA 94956-1346

January 2009

U.S. Department of the Interior  
National Park Service  
Natural Resource Program Center  
Fort Collins, Colorado

The Natural Resource Publication series addresses natural resource topics that are of interest and applicability to a broad readership in the National Park Service and to others in the management of natural resources, including the scientific community, the public, and the NPS conservation and environmental constituencies. Manuscripts are peer-reviewed to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and is designed and published in a professional manner.

Natural Resource Reports are the designated medium for disseminating high priority, current natural resource management information with managerial application. The series targets a general, diverse audience, and may contain NPS policy considerations or address sensitive issues of management applicability. Examples of the diverse array of reports published in this series include vital signs monitoring plans; monitoring protocols; "how to" resource management papers; proceedings of resource management workshops or conferences; annual reports of resource programs or divisions of the Natural Resource Program Center; resource action plans; fact sheets; and regularly-published newsletters.

Views, statements, findings, conclusions, recommendations and data in this report are solely those of the author(s) and do not necessarily reflect views and policies of the U.S. Department of the Interior, NPS. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the National Park Service.

Printed copies of reports in these series may be produced in a limited quantity and they are only available as long as the supply lasts. This report is also available from the Natural Resource Publications Management website (<http://www.nature.nps.gov/publications/NRPM>) and the North Coast and Cascades Network Inventory and Monitoring website (<http://science.nature.nps.gov/im/units/NCCN>) on the Internet, or by sending a request to the address on the back cover.

Please cite this publication as:

Siegel, R. B. 2009. Methods for monitoring landbirds: a review commissioned by Seattle City Light's Wildlife Research Advisory Committee (2000). Natural Resource Report NPS/NCCN/NRR—2009/074. National Park Service, Fort Collins, Colorado.

# Contents

|   |    |
|---|----|
| Acknowledgments.....  | iv |
| Purpose of this report .....  | 1  |
| Long-term natural avian monitoring in the national parks .....              | 1  |
| Review of established landbird monitoring techniques .....                  | 2  |
| Abundance monitoring techniques .....                                       | 3  |
| Area Searches: .....  | 3  |
| Fixed/Unlimited Radius Point Counts without Detectability Estimation: ..... | 4  |
| Point Counts with Detectability Estimation:.....                            | 5  |
| Transect Counts: .....  | 7  |
| Spot Mapping:.....  | 8  |
| Demographic Monitoring Techniques .....                                     | 9  |
| Constant Effort Mist Netting: .....   | 9  |
| Nest Monitoring:.....   | 10 |
| Nest Monitoring with Color Banding: .....                                   | 11 |
| Conclusions.....  | 13 |
| Literature cited.....   | 15 |

## Acknowledgments

I thank Jon Bart, David DeSante, Steve Fancy, Bob Kuntz, Paul Geissler, C. J. Ralph, and John Sauer for helpful discussions of the ideas in this paper. David DeSante, Bob Kuntz, and Phil Nott also provided valuable editorial comments on the manuscript. This is Contribution No. 130 of The Institute for Bird Populations. Photograph of Common Yellowthroat (*Geothlypis trichas*) on the cover was provided by Harold Christenson.

## Purpose of this report

In January 2000 the Wildlife Research Advisory Committee (WRAC) of Seattle City Light awarded North Cascades National Park Service Complex (NOCA) a grant to develop a long-term landbird monitoring plan for North Cascades National Park. The WRAC stipulated that NOCA use part of the grant to review and summarize existing literature on landbird monitoring methods, in order to identify and fully consider the various approaches that might be appropriate for implementation in NOCA and other national parks. This document summarizes our findings.

## Long-term natural avian monitoring in the national parks

Long-term monitoring is increasingly recognized as a crucial ingredient in the effective management of natural resources (Noss and Cooperrider 1994). Nowhere is this more true than within the National Park Service. In 1998 Congress passed the National Parks Omnibus Management Act mandating a “program of inventory and monitoring of National Park System resources to establish baseline information and to provide information on the long-term trends in the condition of National Park system resources.” In fulfillment of this directive, the Servicewide Inventory and Monitoring (I&M) Program of the National Park Service (NPS) has launched a major effort “to encourage, coordinate, standardize, and improve natural resource inventory and monitoring projects throughout the 265 parks with significant natural resources” (Anonymous 1999).

NPS guidelines define natural resource monitoring as “long-term systematic repetition of a specific resource survey and the analysis of those data to predict or detect natural and human-induced changes in resource condition, and to determine if natural resource condition objectives are being achieved”(Anonymous, undated). Numerous authors have outlined why natural resource monitoring is important; one of the most concise, broadly applicable statements comes from Davis (1993):

*“What to monitor, and the appropriate level of accuracy, varies from area to area, but the basic reasons for monitoring are the same everywhere. They are to:*

- Determine present and future health of natural area ecosystems.*
- Establish empirical limits of variation in natural area resources.*
- Diagnose abnormal conditions to identify issues in time to develop effective mitigation, and*
- Identify potential agents of abnormal change.”*

The NPS considers long-term monitoring “necessary to enable managers to make better informed management decisions, to provide early warning of abnormal conditions in time to develop effective mitigation measures, to convince other agencies and individuals to make decisions benefitting parks, to satisfy certain legal mandates, and to provide reference data from relatively pristine areas for comparison with data collected outside of parks by other agencies” (Fancy, undated).

As numerous authors have pointed out, the first step in designing any monitoring program is to define explicit objectives (Silsbee and Peterson 1991). Fancy and Sauer (2000) recently articulated a hierarchy of four general objectives for park managers interested in initiating avian inventory or monitoring projects:

*Objective 1: Document which species occur in the park.*

*Objective 2: Determine distribution and get a qualitative measure of relative abundance (‘abundant’, ‘common’, ‘rare’) of each species in the park.*

*Objective 3: Compare relative abundances among species, habitats or areas, or detect trends in population size.*

*Objective 4: Determine the causes of population trends and differences in abundance among species, habitats, and areas or identify and evaluate management actions to reverse declining trends and increase low population sizes.*

These objectives are presented in increasing order of complexity. Objectives 3 and 4 require a substantially greater investment of resources than do Objectives 1 and 2, but they are also likely to provide managers with much more useful information. Indeed, Objectives 1 and 2 alone are probably not appropriate for meeting the long-term monitoring needs of most parks, as they are unlikely to allow detection of non-catastrophic changes over time, or to facilitate scientifically-based management actions even if catastrophic population declines are detected. Objectives 1 and 2 may, however, be appropriate for shorter-term inventory efforts.

## **Review of established landbird monitoring techniques**

As numerous authors have pointed out, there is no ideal monitoring method; different monitoring techniques are appropriate for satisfying different study objectives. Nevertheless, utilizing standardized methods and protocols can facilitate inter-park comparisons, as well comparisons of park data with data from non-park sources. Unfortunately, such standardization does not



currently exist (Quinn and van Riper III 1990, Sauvajot et al. 1990). Park monitoring efforts thus must strike a careful balance— they must be flexible enough to allow managers to tailor monitoring techniques to particular park needs, while still being sufficiently standardized to allow for intra- and inter-park comparisons, as well as comparisons with data from surrounding lands that may not be under park management.

## **Abundance monitoring techniques**

The following techniques provide data of varying quality and detail on species occurrence, abundance, and/or density: area searches, linear/strip transects, fixed/unlimited radius point counts, point counts with detectability estimation, and spot mapping. All of these methods are limited to varying degrees by a common set of inherent error sources in counting birds. The majority of count detections are aural, at least in densely vegetated habitats, so birds that are not singing may be essentially uncountable. Songbirds show enormous variation in song frequency according to such factors as weather conditions, population density, time of day, and stage of reproduction (Conner and Dickson 1980, Diehl 1981, Mayfield 1981, Robbins 1981, Skirvin 1981). In many species unmated males sing more frequently than mated males (Nolan 1978, Best 1981), and whether they are singing frequently or not, the presence of large numbers of non-breeding ‘floaters’ can greatly skew results (van Riper III 1981). Additionally, the efficiency of even experienced observers decreases significantly as the number of singing birds increases (Bart and Schoultz 1984). For all these reasons, none of the abundance monitoring techniques are perfect. Each technique also has its own additional weaknesses and strengths, which I review and discuss in turn below. I also suggest which of the four previously discussed Objectives can be appropriately pursued using each monitoring technique, and also briefly discuss site selection criteria for implementing each of the methods. Although I don’t explicitly state this in the site selection criteria for each method, survey sites for all bird monitoring techniques should be located to coincide with monitoring locations for studies of plants and other animal taxa whenever feasible, so that habitat relations can be studied as thoroughly and efficiently as possible.

### ***Area Searches:***

The least regimented of the established methods for counting birds, the area search method simply requires that observers roam freely for a fixed time in a specified area, tallying numbers of each species detected (Loyn 1986, Slater 1994). The method is particularly well suited to

projects that rely on volunteer observers (Nur et al. 1999), as it allows variable numbers of observers with diverse skill levels to work together. Because of its informality (observers can behave much as they would during any other day of birding), the method lends itself well to events that combine bird monitoring with public outreach; volunteer observers may wish to participate simply for the fun of it. Furthermore, unlike most other bird-counting techniques, area searches allow rare or secretive species to be actively pursued, perhaps elevating their detection probability (Siegel in preparation).

Area search data also have substantial limitations. Although observers are generally asked to estimate and tally the numbers of individual birds detected, such estimates must be interpreted with great caution. Ensuring that individual birds are counted no more than once can sometimes be difficult even when the observer remains in a fixed location (i.e. during point counts); doing so when multiple observers are moving freely is exponentially harder. Area searches are consequently most appropriate for determining which species are present in a given area, and perhaps qualitatively classifying the relative abundance of each species (rare, common, etc.), but are not appropriate for tracking numbers of birds over time. Even when the objective is simply to generate species lists, area search results may vary substantially with observer skill level and motivation, particularly if the study area includes habitat patches that may be uninviting to some observers (i.e. steep areas, muddy areas, areas dominated by spiny or poisonous plants).

*Suitable objectives:* 1, 2

*Criteria for site selection:* Area searches most effectively survey relatively limited, well-delineated areas (Siegel, in preparation). Survey areas may be circumscribed by man-made boundaries (roads, fences, etc.) or natural boundaries (streams, well-defined ecotones, etc.), but in general, clearer borders around the survey area make for more reproducible results. Similarly, observers surveying relatively small parcels of land that can be thoroughly traversed (perhaps repeatedly) during the search will also provide more reproducible data.

### ***Fixed/Unlimited Radius Point Counts without Detectability Estimation:***

Point counts require observers to record all birds detected during a fixed amount of time (generally between 3 and 10 minutes) within a pre-determined (usually 50 m) and/or unlimited distance of selected survey points (Hutto et al. 1986, Ralph et al. 1993). The resulting data yield abundance indices, under the assumption that the number of birds detected is proportional to the number of birds actually present.

Like transect surveys (see below), point counts are very cost-effective, and can be implemented across large study areas. Point counts have a major advantage over transect surveys, however, in that bird detections are linked to specific locations, where habitat composition and structure, as well as other ecological variables, can be quantified. Additionally, stationary observers may be better able to focus on birds than observers who are walking, and the practice of counting birds while remaining still for several minutes may be less disruptive of normal bird activity than transect counts, where the observer is continuously moving (Reynolds et al. 1980). An additional advantage of point count data is that they can readily be compared with data from the North American Breeding Bird Survey (BBS), which are obtained from roadside point counts (Droege 1990, Peterjohn and Sauer 1993).

The primary drawback of fixed/unlimited radius point counts, as with any survey method that does not take into account detection probability, is that results may vary widely with observer ability and habitat characteristics (Burnham 1981, Barker and Sauer 1995, Fancy and Sauer 2000). The number of birds counted at a sampling station is the product of the number of birds present and the proportion of them the observer detects. Without accounting for differences in the proportion of birds detected due to differing habitats, observers, or other conditions, the appropriateness of pooling point count results among different observers, or comparing results between different habitats, can be called into question (Fancy 1997, Fancy and Sauer 2000).

*Suitable objectives:* 1, 2, 3 (although point counts without detectability estimation have been widely used to pursue objective 3, some authors make a compelling argument that they should not be).

*Criteria for site selection:* Site selection guidelines appropriate for point count studies in national parks are provided by Fancy (2000) and Geissler (undated). Major issues include clear delineation of a sampling frame, incorporation of probabilistic sampling, and targeting extra resources (higher sampling intensity) at areas of special interest.

### ***Point Counts with Detectability Estimation:***

Two distinct methods currently exist for adapting point count methods to account for varying detection probabilities associated with different habitats and observers: distance estimation and the double-observer method. In point counts with distance estimation (often called variable circular plots, or ‘VCPs’), the observer records the distance to each bird detected; analysis of data can then elucidate detection probabilities of birds at increasing distances from the observer

(Buckland et al. 1993). The recently developed double-observer approach requires that two observers simultaneously conduct point counts; detection probabilities are calculated based on the proportion of all birds detected that each individual observer detects (Nichols et al. 2000).

The VCP method yields estimates of detectability, and by extension, meaningful bird density estimates, with only a nominal increase in investment over fixed/unlimited radius point counts without detectability estimation. The double-observer approach may be somewhat more costly to implement, as it requires twice the personnel to survey the same number of points. However, survey protocols in remote wilderness areas often require that observers work in pairs anyway, for safety reasons; in these cases, additional expenses incurred by the double-observer technique may be minimal. In circumstances where personnel is not a limiting factor, the double-observer approach may actually be preferable to the VCP technique, as it avoids errors associated with inaccurate distance estimation (see below). Nevertheless, the double-observer approach has its own set of problems that arise in both field application and data analysis (see Nichols et al. 2000).

VCPs provide substantial advantages over point counts without detectability estimation, and may be somewhat simpler and less costly to implement than double-observer point counts, and have therefore been explicitly endorsed by the National Park Service Inventory and Monitoring Program (Fancy and Sauer 2000). Nevertheless, distance estimation should not be viewed as a panacea for eliminating all important sources of error in conducting point counts. Accuracy of abundance estimates is limited by observer ability to reliably estimate distances; this can be surprisingly difficult, especially in habitats with dense vegetation, high canopy, or rugged topography (DeSante 1986, Hutto et al. 1986, Nur et al. 1999). Relatively modest errors in distance estimation can result in large errors in density estimation (DeSante 1981). Additionally, there may be other detectability issues in bird sampling that neither VCPs nor the double-observer approach can address, such as the existence of unobservable portions of the population being surveyed (e.g. highly secretive females)(Fancy and Sauer 2000).

*Suitable objectives:* 1, 2, 3

*Criteria for site selection:* See criteria for fixed/unlimited radius point counts without detectability estimation.

### ***Transect Counts:***

Transect counts include a diverse set of protocols with important distinctions between them. Terminology for the different protocols is quite variable in the published literature, but Verner (1985) provides a thorough clarification and review. Verner (1985) and Franzreb (1981) provide extensive discussions of the assumptions implicit in each of the major versions of the transect method.

Line transects (also called variable-width transects) and strip transects (also called fixed-width transects) require an observer to record all birds detected while walking a fixed route (Emlen 1971). Most line transect protocols require the observer to estimate the perpendicular distance to each bird detected, and consequently allow the calculation of detection probability as a function of distance from the transect. Strip transects require the observer to record all birds detected within a fixed distance of the transect (often 50m), and sometimes additionally require that all birds beyond 50m be recorded separately.

It has been argued that transects provide a more efficient means of counting birds than do point counts (Anderson and Ohmart 1981, Bibby et al. 1992, Fancy and Sauer 2000) as they allow observers to spend more of their field time actually counting birds; a greater effective survey area can be covered, yielding higher detection rates, and consequently, greater precision. On the other hand, overall statistical power of the method is weakened by the fact that it yields fewer independent data points or replicates than point count methods (Nur et al. 1999). Additionally, the method is generally feasible only on relatively gentle terrain, where observers are able to focus their attention fully on birds, rather than on watching their own footing (Anderson and Ohmart 1981, Dawson 1981). Finally, the largest shortcoming of the transect count methods is that they fail to yield associations between individual bird detections and particular points in space. Since the specific locations of individual detections along a transect are generally not recorded, potentially valuable information about habitat associations is lost (Fancy and Sauer 2000). This is particularly true when transects pass through varied habitats. Transect methods are therefore not appropriate for surveying birds in patchy or fine-grained habitats (Anderson and Ohmart 1981, Bibby et al. 1992), or for studies that seek to correlate relatively fine-scale habitat variables with bird occurrence.

*Suitable objectives:* 1, 2, 3 (if detection probability is incorporated)

*Criteria for site selection:* If study goals include clarifying bird-habitat relationships, individual

transects must be located wholly within a single habitat type.

***Spot Mapping:***

First described in the literature by Kendeigh (1944), spot mapping as a monitoring technique was later explicitly endorsed by the International Bird Census Committee (International Bird Census Committee 1969), which provided guidelines for its use (Anonymous 1970, Robbins 1970). Although the details of the method have been modified over the years (Svensson 1979 and 1980, Tomialojc 1980) the basic technique is to mark locations, movements, and interactions of individual birds on detailed maps of the study area. Information gathered from multiple visits can be used to map individual breeding territories within the study area.

The spot mapping technique has several distinct strengths. Unlike abundance monitoring techniques that do not incorporate detectability estimation, spot mapping provides density estimates, rather than merely indices of abundance. Additionally, because the technique estimates density of territorial birds, the results may be more meaningful than density estimates produced by point counts with detectability estimation, as the point count estimates may be biased high due to the presence of nonbreeding floaters. Finally, the method can potentially produce more detailed natural history information (i.e. habitat usage, breeding behavior, etc.) than most other monitoring techniques.

Spot mapping also has several important limitations. Because the technique involves multiple visits to the study site, it can be quite expensive per data point (Bibby et al. 1992), a factor that led Nur et al. (1999) to conclude, “[spot mapping] may be better applied to research projects or to high priority areas or species.” Additionally, the technique can generally be implemented only in a fairly circumscribed study area, and can therefore be of only limited value if researchers wish to extrapolate results across a large park. Some authors have also cautioned that inferring territory boundaries from the movements and territorial interactions of individual birds can be highly subjective, with resulting density estimates varying widely among observers (Oelke 1981, Verner 1985). Finally, spot mapping is the only abundance monitoring technique discussed here that is applicable solely during the breeding season.

*Suitable objectives:* 3

*Criteria for site selection:* Spot mapping studies are usually limited to only a small number of relatively small study plots, because of the substantial time investment necessary to complete the mapping. To provide maximum potential for extrapolating results to the surrounding area, study

plots 1) should contain habitat that is representative of the larger landscape, and 2) should be sited using a probabilistic sampling strategy. In some landscapes, satisfying both these conditions can be quite challenging.

## **Demographic Monitoring Techniques**

In addition to measuring avian abundance, the following methods also estimate one or more demographic parameters, such as nesting productivity, post-fledging productivity, recruitment, or survivorship. Monitoring demographic parameters is a crucial component of meaningful landbird monitoring efforts for several reasons. First, environmental stressors and management actions affect demographic parameters directly, without the buffering or time lags that so often affect secondary population trends (Temple and Wiens 1989). Because of the buffering effects of floater individuals and the density-dependent responses of populations, there may be substantial time lags between changes in primary demographic parameters and resulting changes in population size or density (DeSante and George 1994). Second, demographic parameters elucidate the stage(s) in the life cycle at which population change is being affected (DeSante 1992). Such information is particularly important for migratory birds, as it can determine whether management actions should be directed toward a species' breeding grounds, wintering grounds, or both. Finally, monitoring demographic parameters provides information critical for assessing the viability of populations, and for identifying areas of source-sink population dynamics (Donovan et al. 1995). Indeed, because of the vagility of most bird species, local variation in reproductive success may often be masked by recruitment from a wider region (George et al. 1992) or accentuated by lack of recruitment from a larger area (DeSante 1990). Source-sink dynamics may thus make the density of a species in a given area a misleading indicator of population viability (Van Horne 1983, Pulliam 1988).

### ***Constant Effort Mist Netting:***

Constant effort mist netting can provide indices of post-fledging productivity, and estimates of adult population size, adult survivorship, and recruitment into the adult population for target species (Baillie 1990, DeSante et al. 1995, Peach et al. 1996, Bart et al. 1999, Silkey et al. 1999). The widespread use of the Monitoring Avian Productivity and Survivorship (MAPS) constant effort mist netting protocol across North America (DeSante et al. 1998, DeSante et al. 2000) allows parameter values to be compared or combined with mist netting results from other areas at multiple spatial scales. The mark-recapture techniques incorporated by MAPS provide the

only methodology (other than resighting individually color marked birds-- see nest monitoring with color banding, below) that permits estimation of adult survivorship and recruitment into the adult population. These parameters are critical for providing a full understanding of demography and population dynamics. Additionally, constant effort mist netting allows numerous species to be monitored at the same time, without additional effort, and in-hand examination of birds facilitates collection of additional natural history data.

Like other demographic monitoring techniques, mist netting is a relatively intensive, rather than extensive approach to monitoring, so the spatial extent of the area that can be effectively surveyed is limited. A single MAPS station, with an array of ten 12m mist nets, covers an area of approximately 20-ha, although productivity indices reflect reproductive success from the larger landscape (DeSante et al. 2000). At least four consecutive years of data collection are necessary to produce survivorship estimates for locally resident adult birds, but productivity estimates may be obtained from a single year of data collection. As a general rule of thumb, an average of at least seven individual adult birds per year of each target species must be captured to produce robust survivorship estimates (D. DeSante, pers. comm.). These sample size requirements for parameter estimation makes the technique impractical for monitoring locally rare species. Finally, mist netting is an ineffective monitoring technique for bird species whose habits restrict them to grasslands or to the upper forest canopy.

*Suitable objectives:* 3, 4

*Criteria for site selection:* DeSante et al. (2000) provide detailed recommendations for siting constant effort mist netting stations. In general, they suggest that stations be sited in habitat types that are fairly representative of the surrounding landscape, that stations include some edge habitat (such as forest edge, riparian corridor, montane meadow, or power-line right-of-way) to help ensure that large numbers of bird will be captured, and that, to the extent possible, station locations are chosen under a probabilistic sampling strategy.

### ***Nest Monitoring:***

Detailed field protocols and pointers for finding and monitoring nests are provided in Martin and Geupel (1993) and Martin et al. (1997). Searching for nests of target species and tracking nesting attempts throughout the breeding season is the only way to effectively estimate nest success rates and to determine proximate causes of nest failures. Nest monitoring facilitates the study of nest-sites or characteristics, and by extension, the elucidation of correlations between nest sites or



habitat characteristics and nest success or failure (Best and Stauffer 1980, Li and Martin 1991, Martin 1993, Robinson et al. 1995).

Like mist netting, the spatial extent and habitat diversity of the area that can be effectively surveyed through nest monitoring is somewhat limited. Unlike mist netting, which generally provides post-fledging productivity estimates applicable to the larger landscape (because many of the hatching-year birds caught are dispersing, rather than locally produced individuals), nest monitoring yields fledging success rates that are specific to the study plot— this can be an advantage or a disadvantage, depending on study objectives. Unless all nesting attempts in the study area are monitored, however, the actual reproductive output of the various species cannot be determined. Moreover, nest monitoring is quite labor-intensive, generally requiring that nests be visited at least every 3-4 days (more frequently is even better) to allow for reliable determinations of nest fates (Martin et al. 1997). For this reason the method is generally better suited for testing specific management-related hypotheses than for long-term monitoring in the strict sense.

An additional drawback to nest monitoring is that many studies do not attain large enough sample sizes for robust statistical comparisons. Hensler and Nichols (1981) provide a table of sample sizes necessary to estimate nest survivorship under different parameter values, and conclude that data from a minimum of 20 nests must be pooled to obtain nest survivorship estimates with acceptable precision. Nur et al. (1999) extend the analysis to look at sample size requirements for providing enough statistical power to compare nest survivorship rates among treatments, and conclude that 75 nests per group is a more reliable rule of thumb.

*Suitable objectives:* 3, 4

*Criteria for site selection:* Detailed guidelines for siting nest monitoring plots are provided by Martin et al. (1997). As with siting mist netting stations, researchers should use a randomization scheme to randomly choose study plots from all possible sites meeting pre-determined ecological or logistical criteria.

### ***Nest Monitoring with Color Banding:***

Referred to as ‘total mapping’ by Verner (1985), nest monitoring combined with target netting and color banding of breeding individuals and nestlings, and systematic efforts to resight color banded birds is the most thorough means of monitoring individual bird populations. If all nesting

attempts are monitored and the fates of fledglings are followed to independence from their parents, this suite of techniques provides the most complete and unbiased measures of demographic parameters of any method (Nur et al. 1999). This method, however, is also extremely labor-intensive, much more so than any other method discussed in this review. For this reason the method is not appropriate for general monitoring programs, but should instead be reserved only for high-priority species or areas (Nur et al. 1999).

*Suitable objectives:* 3, 4

*Criteria for site selection:* Site selection criteria for monitoring projects employing nest monitoring and color banding will usually be driven by the particular circumstances requiring such a labor-intensive mode of study.

## Conclusions

Table 1 summarizes which Objectives can potentially be attained through each monitoring method discussed in this review. It should be noted, however, that the different methods yield data of substantially different nature and quality, even methods deemed appropriate for the same Objectives. Table 2 provides a summary of the kinds of data yielded by each survey method.

Researchers designing long-term monitoring projects are likely to conclude that no single monitoring method will fully meet their needs; abundance monitoring techniques are relatively inexpensive and can be implemented across very large areas, but provide little information about the causes of population changes that may be detected. In contrast, demographic monitoring techniques can give valuable insight into such causes, but are generally much more expensive per datum, and can only be implemented across relatively small study plots. For these reasons, many authors assert that monitoring programs should ideally incorporate elements of both abundance monitoring and demographic monitoring techniques (Taylor et al. 1985, Verner 1985, Baillie 1990, Nur and Geupel 1993, Ralph et al. 1993, DeSante 1995, Saab and Rich 1997, DeSante and Rosenberg 1998). Integrating spatially extensive abundance surveys with more intensive demographic studies of carefully selected target species or representative locations is likely to yield data of maximum use to park managers interested in both tracking landbird populations over time and implementing appropriate remedial management actions should it become clear that populations are threatened.



## Literature cited

- Anderson, B. W. and R. D. Ohmart. 1981. Comparison of avian census results using variable distance transect and variable circular plot techniques. Pp. 186-192 *in* C. J. Ralph and M. Scott (eds.), Estimating numbers of terrestrial birds, Studies in Avian Biology No. 6, Cooper Ornithological Society.
- Anonymous. Undated. Natural resources inventory and monitoring guideline. NPS-75. National Park Service Inventory and Monitoring Program.
- Anonymous. 1970. Recommendations for an international standard for a mapping method in bird census work. Swedish Nat. Sci. Res. Council, Stockholm. Bull. Ecol. Res. Comm. 9:49-52.
- Anonymous. 1999. Guidelines for biological inventories. Inventory and Monitoring Program, National Park Service.
- Baillie, S. R. 1990. Integrated population monitoring of breeding birds in Britain and Ireland. *Ibis* 132:151-166.
- Barker, R. J. and J. R. Sauer. 1995. Statistical aspects of point count sampling. Pp. 125-130 *in* C.J. Ralph, J.R. Sauer and S. Droege (eds.), Monitoring bird populations by point counts, USDA Forest Service, Pacific Southwest Research Station, Gen. Tech. Rep. PSW-GTR-149.
- Bart, J., C. Kepler, P. Sykes, and C. Bocetti. 1999. Evaluation of mist-net sampling as an index to productivity in Kirtland's Warblers. *Auk* 116:1147-1151.
- Bart, J. and J. D. Schoultz. 1984. Reliability of singing birds surveys: changes in observer efficiency with avian density. *Auk* 101:307-318.
- Best, L. B. 1981. Seasonal changes in detection of individual bird species. Pp. 252-261 *in* C. J. Ralph and M. Scott (eds.), Estimating numbers of terrestrial birds, Studies in Avian Biology No. 6, Cooper Ornithological Society.

- Best, L. B. and D. F. Stauffer. 1980. Factors affecting nesting success in riparian bird communities. *Condor* 82:149-158.
- Bibby, C. J., N. D. Burgess, and D. A. Hill. 1992. *Bird census techniques*. Academic Press, London.
- Buckland, S. T., D. R. Anderson, K. P. Burnham, and J. L. Laake. 1993. *Distance sampling: estimating abundance of biological populations*. Chapman and Hall, New York.
- Burnham, K. P. 1981. Summarizing remarks: environmental influences. *Studies in Avian Biology* 6:324-325.
- Conner, R. N. and J. G. Dickson. 1980. Strip transect sampling and analysis for avian habitat studies. *Wildl. Soc. Bull.* 8:4-10.
- Davis, G. E. 1993. Design elements of monitoring programs: the necessary ingredients for success. *Environmental Monitoring and Assessment* 26:99-105.
- Dawson, D. K. 1981. Sampling in rugged terrain. Pp. 311-315 *in* C. J. Ralph and M. Scott (eds.), *Estimating numbers of terrestrial birds*, *Studies in Avian Biology* No. 6, Cooper Ornithological Society.
- DeSante, D.F. 1981. A field test of the variable circular-plot censusing technique in a California coastal scrub breeding bird community. Pp. 177-185 *in* C. J. Ralph and M. Scott (eds.), *Estimating numbers of terrestrial birds*, *Studies in Avian Biology* No. 6, Cooper Ornithological Society.
- DeSante, D. F. 1986. A field test of the variable circular-plot censusing method in a Sierran subalpine forest habitat. *Condor* 88:129-142.
- DeSante, D. F. 1990. The role of recruitment in the dynamics of a Sierran subalpine bird community. *American Naturalist* 136:429-455.

- DeSante, D. F. 1992. Monitoring Avian Productivity and Survivorship (MAPS): A sharp, rather than blunt, tool for monitoring and assessing landbird populations. Pp. 511-521 *in* D. R. McCullough and R. H. Barrett (eds.), *Wildlife 2001: Populations*. Elsevier Applied Science, London, UK.
- DeSante, D. F. 1995. Suggestions for future directions for studies of marked migratory landbirds from the perspective of a practitioner in population management and conservation. *Journal of Applied Statistics* 22:944-965.
- DeSante, D. F., K. Burton, J. F. Saracco, and B. L. Walker. 1995. Productivity indices and survival rate estimates from MAPS, a continent-wide programme of constant-effort mist netting in North America. *Journal of Applied Statistics* 22:935-947.
- DeSante, D. F., K. Burton, P. Velez and D. Froehlich. 2000. MAPS manual. The Institute for Bird Populations, Point Reyes Station, CA.
- DeSante, D. F. and T. L. George. 1994. Population trends in the landbirds of western North America. Pp. 173-190 *in* J. R. Jehl, Jr. and N. K. Johnson (eds.), *A century of avifaunal change in North America*, Studies in Avian Biology No 15, Cooper Ornithological Society.
- DeSante, D. F., D. R. O'Grady, K. M. Burton, P. Velez, D. Froehlich, E. E. Feuss, H. Smith and E. D. Ruhlen. 1998. The Monitoring Avian Productivity and Survivorship (MAPS) Program sixth and seventh annual report (1995 and 1996). *Bird Populations* 4:69-122.
- DeSante, D. F. and D. K. Rosenberg. 1998. What do we need to monitor in order to manage landbirds? Pp. 93-106 *in* J. M. Marzluff and R. Sallabanks (eds.), *Avian conservation: research and management*. Island Press, Washington, D.C.
- Diehl, B. 1981. Bird populations consist of individuals differing in many respects. Pp. 225-229 *in* C. J. Ralph and M. Scott (eds.), *Estimating numbers of terrestrial birds*, Studies in Avian Biology No. 6, Cooper Ornithological Society.
- Donovan, T. M., F. R. Thompson III, J. Faaborg, and J. R. Probst. 1995. Reproductive success of migratory birds in habitat sources and sinks. *Conservation Biology* 9:1380-1395.

- Droege, S. 1990. The North American Breeding Bird Survey. Pp. 1-4 *in* J. R. Sauer and S. Droege(eds.), Survey design and statistical methods for the estimation of avian population trends. U.S. fish and Wildlife Service, Biol. Rep. 90(1).
- Emlen, J. T. 1971. Population densities of birds derived from transect counts. *Auk* 88:323-342.
- Fancy, S. G. Undated. Monitoring natural resources in our national parks. National Park Service Inventory and Monitoring Program.
- Fancy, S. G. 1997. A new approach for analyzing bird densities from variable circular-plot counts. *Pacific Science* 51:107-114.
- Fancy, S. G. 2000. Guidance for the design of sampling schemes for inventory and monitoring of biological resources in national parks. National Park Service Inventory and Monitoring Program.
- Fancy, S. G. and J. R. Sauer. 2000. Recommended methods for inventorying and monitoring landbirds in national parks. National Park Service Inventory and Monitoring Program.
- Franzreb, K. E. 1981. The determination of avian densities using the variable-strip and fixed-width transect surveying methods. Pp. 139-145 *in* C. J. Ralph and M. Scott (eds.), Estimating numbers of terrestrial birds, Studies in Avian Biology No. 6, Cooper Ornithological Society.
- Geissler, P. Undated. National park inventories: statistical methods. USGS Biological Resources Division.
- George, T. L., A. C. Fowler, R. L. Knight, and L. C. McEwen. 1992. Impacts of a severe drought on grassland birds in western North America. *Ecological Applications* 2:275-284.
- Hensler, G. L. and J. D. Nichols. 1981. The Mayfield method of estimating nesting success: a model, estimators and simulation results. *Wilson Bull.* 93:42-53.
- Hutto, R. L., S. M. Pletschet, and P. Hendricks. 1986. A fixed-radius point count method for nonbreeding and breeding season use. *Auk* 103:593-602.



- International Bird Census Committee. 1969. Recommendations for an international standard for a mapping method in bird census work. *Bird Study* 16:248-255.
- Kendeigh, S. C. 1944. Measurement of bird populations. *Ecol. Monogr.* 14:67-106.
- Li, P. and T. T. Martin. 1991. Nest-site selection and nesting success of cavity-nesting birds in high elevation forest drainages. *Auk* 108:405-418.
- Loyn, R. H. 1986. The 20 minute search— a simple method for counting forest birds. *Corella* 10:58-60.
- Martin, T. E. 1993. Nest predation and nest sites. *Bioscience* 43:523-532.
- Martin, T. E. and G. R. Geupel. 1993. Nest monitoring plots: methods for locating nests and monitoring success. *J. Field Ornithol.* 64:507-519.
- Martin, T. E., C. Paine, C. J. Conway, W. M. Hochachka, P. Allen, and W. Jenkins. 1997. BBIRD field protocol. Montana Cooperative Wildlife Research Unit, University of Montana, Missoula, MT.
- Mayfield, H. F. 1981. Problems in estimating population size through counts of singing males. Pp. 220-224 *in* C. J. Ralph and M. Scott (eds.), *Estimating numbers of terrestrial birds*, Studies in Avian Biology No. 6, Cooper Ornithological Society.
- Nichols, J. D., J. E. Hines, J. R. Sauer, F. W. Fallon, J. E. Fallon, and P. J. Heglund. 2000. A double-observer approach for estimating detection probability and abundance from point counts. *Auk* 117:393-408.
- Nolan, V. 1978. The ecology and behavior of the Prairie Warbler, *Dendroica discolor*. *Ornithol. Monogr.* 26.
- Noss, R. F. and A. Y. Cooperrider. 1994. *Saving nature's legacy: protecting and restoring biodiversity*. Island Press, Washington, D.C.

- Nur, N. and G. R. Geupel. 1993. Evaluation of mist-netting, nest-searching and other methods for monitoring demographic processes in landbird populations. Pp. 237-244 *in* D. M. Finch and P. W. Stangel (eds.), Status and management of Neotropical migratory birds. Gen. Tech. Rep. RM-229, USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- Nur, N, S. L. Jones, and G. R. Guepel. 1999. A statistical guide to data analysis of avian monitoring programs. U.S. Department of the Interior, Fish and Wildlife Service, BTP-R6001-1999, Washington, D.C.
- Oelke, H. 1981. Limitations of the mapping method. Pp. 114-118 *in* C. J. Ralph and M. Scott (eds.), Estimating numbers of terrestrial birds, Studies in Avian Biology No. 6, Cooper Ornithological Society.
- Peach, W. J., S. T. Buckland, and S. R. Baillie. 1996. The use of constant effort mist-netting to measure between-year changes in the abundance and productivity of common passerines. *Bird Study* 43:142-156.
- Peterjohn, B.G. and J. R. Sauer. 1993. North American breeding bird survey annual summary 1990-1991. *Bird Populations* 1:1-15.
- Pulliam, H. R. 1988. Sources, sinks, and population regulation. *American Naturalist* 132:652-661.
- Quinn, J. F. and van Riper III, C. 1990. Design considerations for national park inventory databases. Pp. 5-13 *in* C. van Riper III, T. J. Stohlgren, S D. Veirs, and S. Castillo Hillyer (eds.), Examples of resource inventory and monitoring in national parks of California. U.S. Department of the Interior, National Park Service, Washington, D.C.
- Ralph, C. J., G. R. Geupel, P. Pyle, T. E. Martin, D. F. DeSante. 1993. Handbook of field methods for monitoring landbirds. Gen. Tech. Rep. PSW-GTR-144. Albany, CA: Pacific Southwest Research Station, Forest Service, USDA.
- Reynolds, R. T., J. M. Scott, and R. A. Nussbaum. 1980. A variable circular-plot method for estimating bird numbers. *Condor* 82:309-313.

- Robbins, C. S. 1970. Recommendations for an international standard for a mapping method in bird census work. *Aud. Field Notes* 24:723-726.
- Robbins, C. S. 1981. Effect of time of day on bird activity. Pp. 275-286 *in* C. J. Ralph and M. Scott (eds.), *Estimating numbers of terrestrial birds*, *Studies in Avian Biology* No. 6, Cooper Ornithological Society.
- Robinson, S. K., F. R. Thompson III, T. M. Donovan, D. R. Whitehead, and J. Faaborg. 1995. Regional forest fragmentation and the nesting success of migratory birds. *Science* 267:1987-1990.
- Saab, V. A. and T. D. Rich. 1997. Large-scale conservation assessment for neotropical migratory landbirds in the Columbia River Basin. Gen. Tech. Rep. PNW-GTR-399. Pacific Northwest Research Station, USDA Forest Service, Portland, OR.
- Sauvajot, R. M., J. F. Quinn, C. van Riper III, and C. Farmer. 1990. Comparative analyses of bird inventory databases from California national parks. Pp. 15-29 *in* C. van Riper III, T. J. Stohlgren, S D. Veirs, and S. Castillo Hillyer (eds.), *Examples of resource inventory and monitoring in national parks of California*. U.S. Department of the Interior, National Park Service, Washington, D.C.
- Siegel, R. B. In preparation. Comparing the results of area searches and point counts for surveying meadow bird communities.
- Silkey, M., N. Nur, and G. R. Guepel. 1999. The use of mist-net capture rates to monitor annual variation in abundance: a validation study. *Condor* 101:288-298.
- Silsbee, D. G. and D. L. Peterson. 1991. Designing and implementing comprehensive long-term inventory and monitoring programs for national park system lands. *Natural Resources Report NPS/NRUW/NRR-91/04*, National Park Service.
- Skirvin, A. A. 1981. Effect of time of day and time of season on the number of observations and density estimates of breeding birds. Pp. 271-274 *in* C. J. Ralph and M. Scott (eds.), *Estimating numbers of terrestrial birds*, *Studies in Avian Biology* No. 6, Cooper Ornithological Society.

- Slater, P. J. 1994. Factors affecting the efficiency of the area search method of censusing birds in open forests and woodlands. *Emu* 94:9-16.
- Svensson, S. 1979. Census efficiency and number of visits to a study plot when estimating bird densities by the territory mapping method. *Acta Ornithol.* 14:322-338.
- Taylor, K., R. J. Fuller, and P. C. Lack (eds.). 1985. Proceedings of the VIII international conference on bird census and atlas work. *Bird Census and Atlas Studies*, British Trust for Ornithology, Tring, England.
- Temple, S. A. and J. A. Wiens. 1989. Bird populations and environmental changes: can birds be bio-indicators? *American Birds* 43:260-270.
- Tomialojc, L. 1980. The combined version of the mapping method. Pp. 92-106 *in* H. Oelke (ed.), *Bird census work and nature conservation, Proc. VI Intern. Conf. on Bird Census Work*. Univ. Göttingen, Germany.
- Van Horne, B. 1983. Density as a misleading indicator of habitat quality. *Journal of Wildlife Management* 47:893-901.
- van Riper III, C. 1981. Summarizing remarks: comparison of methods. Pp. 217-218 *in* C. J. Ralph and M. Scott (eds.), *Estimating numbers of terrestrial birds*, *Studies in Avian Biology* No. 6, Cooper Ornithological Society.
- Verner, J. 1985. Assessment of counting techniques. *Current Ornithology* 2:247-302

The Department of the Interior protects and manages the nation's natural resources and cultural heritage; provides scientific and other information about those resources; and honors its special responsibilities to American Indians, Alaska Natives, and affiliated Island Communities.

NPS D-235, January 2009

**National Park Service**  
**U.S. Department of the Interior**



---

**Natural Resource Program Center**  
1201 Oakridge Drive, Suite 150  
Fort Collins, CO 80525

[www.nature.nps.gov](http://www.nature.nps.gov)

**EXPERIENCE YOUR AMERICA™**