### Separating Blue-headed and Cassin's Vireos



The Institute for Bird Populations



#### THE INSTITUTE FOR BIRD POPULATIONS

P.O. Box 1346

Point Reyes Station, CA 94956-1346

(415) 663-1436 • FAX (415) 663-9482 • www.birdpop.org

Comments on winter 2010-2011 Ensenada and Point Loma Blue-headed Vireos

Peter Pyle




The Institute for Bird Populations



Dullest HY/SY female BHVI (top, MVZ 43406, SC, 26 Sep 1895) and brightest AHY/ASY male CAVI (MVZ 87287, Tulare County, CA, 30 Aug 1943) at MVZ. Head color in the overlap area is dull blue, breast is medium-bright white, and flanks are medium-dull yellow.









The Institute for Bird Populations

Thus, once age and sex is determined, the two species are more-easily separated.

It is best to age birds first, then assess sex and species simultaneously.

Crisp digital images allow successful ageing and species diagnosis

First-year

#### Blue-headed vs. Cassin's Vireos




The Institute for Bird Populations

Primary coverts brown, contrasting with greater coverts

Primary coverts dusky, not contrasting with greater coverts

Adult Block pattern





The Institute for Bird Populations

Fresh fall



Worn spring

First-year Male BHVI





The Institute for Bird Populations

**Adult Male CAVI** 




# Blue-headed vs. Cassin's Vireos Adult Female BHVI





The Institute for Bird Populations

First-year Male BHVI










# Blue-headed vs. Cassin's Vireos First-year Male BHVI





Adult Male CAVI





First-year Male



